

QEEG Clinical Report

EEGLens



The QEEG report is provided by NPCindex Company, operating under the QEEGhome brand.

Personal Data:

Name: Radin Mahdavi

Gender: Male

Age: 2007-12-04 - 18.1

Handedness: Right

Clinical Data:




Initial diagnosis: R/O ADHD

Medication: -

Date of Recording: 2025-10-08

Source of Referral: Maryam Mirzajanzadeh

This case belongs to **Maryam Mirzajanzadeh**


EEG Quality**Z-score Information****TMS Reponsibility****Pathological Assessment****EEG Neuromarker Values**

Neuromarker	Region	Value	Assessment
AFP	Frontal	11.17	High
AFP	Occipital	11.25	High
Arousal Level	-	-	Normal

Denoising Information

■ Eye Close

Raw EEG



Rejected Channel



Total Recording Time Remaining:

478.57 sec

Number of Eye and Muscle Elements

Eye: 1

Muscle: 0

Low Artifact Percentage



High Artifact Percentage




Total Artifact Percentage



EEG Quality: good

Denoised EEG




Flat Channel



Pathological Assessment

Main Diagnosis: Adult ADHD



Description

According to the guidelines, the initial diagnosis of adult ADHD could have comorbidities such as **alcohol abuse, anxiety, and depression**. It also differentially diagnoses with **depression, anxiety, and BMD**.

In the above graph, the **red area** shows the percentage of each comorbidity from your patient's EEG markers. Observe that each comorbidity marker is not unique and can be shared with other comorbidities.

Side circles in the above graph represent the differential diagnosis between depression and its misdiagnosis conditions based on your patient's EEG markers and trained artificial intelligence. The differential diagnosis probability is represented by **the bold blue bars** in the circles, and the probability of depression is represented by the **gray bars**.


Note: In case your patient has drug abuse, obtain the substance abuse pathologic page of QEEGhome by registering the diagnosis under the initial diagnoses section of the website.

References:

Sadock, B. J., Sadock, V. A., & Ruiz, P. (Eds.). (2025). Kaplan and Sadock's comprehensive textbook of psychiatry (11th ed., Vols. 1–2). Wolters Kluwer
 Sadock, B. J., Sadock, V. A., & Ruiz, P. (2022). Kaplan and Sadock's synopsis of psychiatry: Behavioral sciences/clinical psychiatry (12th ed.). Wolters Kluwer


User Manual

Differential Diagnosis Probability





Comorbidity Percentage

Main Diagnosis Compatibility



QEEG Based Predicting Medication Response



Explanation

These two tables can be considered the most important finding that can be extracted from QEEG. To prepare this list, the NPCIndex Article Review Team has studied, categorized, and extracted algorithms from many authoritative published articles on predict medication response and Pharmacoe EEG studies. These articles are published between 1970 and 2021. The findings extracted from this set include 85 different factors in the raw band domains, spectrum, power, coherence, and loreta that have not been segregated to avoid complexity, and their results are shown in these diagrams. One can review details in NPCIndex.com .

Medication Recommendation

These two charts, calculate response probability to various medications, according only to QEEG indicators. Blue charts favor drug response and red charts favor drug resistance. The longer the bar, the more evidence there is in the articles. Only drugs listed in the articles are listed. These tables present the indicators reviewed in the QEEG studies and are not a substitute for physician selection.

rTMS Response Prediction

Network Performance


Accuracy: 92.10%
Sensitivity: 89.13%
Specificity: 97.47%

Participants Information


Distribution of Gender




Distribution of Age




Features Information



Responsibility




Data Distribution



About Predicting rTMS Response

This index was obtained based on machine learning approaches and by examining the QEEG biomarkers of more than 470 cases treated with rTMS. The cases were diagnosed with depression (with and without comorbidity) and all were medication free. By examining more than 40 biomarkers capable of predicting response to rTMS treatment in previous studies and with data analysis, finally 10 biomarkers including bispectral and nonlinear features entered the machine learning process. The final chart can distinguish between RTMS responsive and resistant cases with 92.1% accuracy. This difference rate is much higher than the average response to treatment of 44%, in the selection of patients with clinical criteria, and is an important finding in the direction of personalized treatment for rTMS.


Vigilance



Vigilance Slope

-0.16


2min



EEG Neuromarker Values

Neuromarker	Region	Value	Assessment
APF	Frontal	11.17	High
APF	Occipital	11.25	High
Alpha Asymmetry	Frontal	-0.15	Anhedonia
Alpha Asymmetry	Occipital	-0.10	Anhedonia
Beta Asymmetry	Frontal	-0.03	Anxiety
Arousal Level	-	-	Normal
Vigilance Level	-	04.00	Normal
Vigilance Mean	-	03.78	Normal
Vigilance Regulation	-	-0.16	Normal
Vigilance 0 Stage (%)	-	00.00	Normal
Vigilance A1 Stage (%)	-	21.55	-

EEG Spectra



Z Score Summary Information


■ Eye Close

Absolute Power

Relative Power


Generation Source

Coherence




Theta/Beta Ratio

■ Eye Close



Absolute Power-Eye Close



Relative Power-Eye Close

