



QEEG Clinical Report

BrainLens V0.4

Report Description



Personal & Clinical Data

Name	Tabasom Gholam Pasand	Date of Recording	17-Nov-2024
Date of Birth - Age	13-Sep-2011 - 13.18	Gender	Female
Handedness(R/L)	Right	Source of Referral	Dr Moradkhani
Initial Diagnosis	Anxiety-Paranoid		
Current Medication	-		


Dr Moradkhani

Summary Report


EEG Quality




Z-score Information



Compatibility with ADHD




Arousal Level



APF

Posterior APF-EC= 11.13


Posterior APF-EO= 11.13




To investigate QEEG-based predicting medication response, please refer to the Report.

Denoising Information (EC)

Raw EEG



Denoised EEG



Flat Channels




Rejected Channels




Number of Eye and Muscle Elements				Low Artifact Percentage	
Eye	1	Muscle	0		
Total Artifact Percentage				High Artifact Percentage	
EEG Quality		bad		Total Recording Time Remaining	
				186.14 sec	

Denoising Information (EO)

Raw EEG



Denoised EEG



Flat Channels




Rejected Channels





Number of Eye and Muscle Elements				Low Artifact Percentage	
Eye	1	Muscle	0		
Total Artifact Percentage				High Artifact Percentage	
EEG Quality		bad		Total Recording Time Remaining	
				182.82 sec	

Pathological assessment for ADHD


Compare to ADHD Database



EEG Compatibility with ADHD Diagnosis



Arousal Level Detection




ADHD Clustering *

1. Prone to moody behavior and temper tantrums. May be anxious, may be highly intelligent, need sufficient sleep, and should avoid high carbohydrate intake. Avoid stimulants, benzodiazepines, SSRI and SNRI. Consider clonidine.
2. Same inattentive and hyperactive prevalence, may be anxious, may be highly intelligent, need sufficient sleep, and should avoid high carbohydrate intake. Consider clonidine

* If there is Paroxymal epileptic discharge in EEG data, this case needs sufficient sleep and should avoid high carbohydrate intake.
You can consider anticonvulant medications.

QEEG based predicting medication response




Explanation

These two tables can be considered the most important finding that can be extracted from QEEG. To prepare this list, the NPCIndex Article Review Team has studied, categorized, and extracted algorithms from many authoritative published articles on predict medication response and Pharmacoe EEG studies. These articles are published between 1970 and 2021. The findings extracted from this set include 85 different factors in the raw band domains, spectrum, power, coherence, and loreta that have not been segregated to avoid complexity, and their results are shown in these diagrams. One can review details in NPCIndex.com .

Medication Recommendation

These two charts, calculate response probability to various medications, according only to QEEG indicators. Blue charts favor drug response and red charts favor drug resistance. The longer the bar, the more evidence there is in the articles. Only drugs listed in the articles are listed. These tables present the indicators reviewed in the QEEG studies and are not a substitute for physician selection.


APF(EO)



Frontal APF= 11.00

Posterior APF= 11.13




APF(EC)




Frontal APF= 11.08

Posterior APF= 11.13


EEG Spectra




Alpha Asymmetry(AA)




Alpha Blocking




Z Score Summary Information (EC)




Z Score Summary Information (EO)





E.C.T/B Ratio (Raw- Z Score)




E.O.T/B Ratio (Raw- Z Score)




Arousal Level




Absolute Power-Eye Closed (EC)



Relative Power-Eye Closed (EC)



Absolute Power-Eye Open (EO)



Relative Power-Eye Open (EO)

